

Carbon-14 Measurements in Atmospheric CO_2 from Northern and Southern Hemisphere Sites, 1962-1993



Reidar Nydal and Knut Lövseth
~ Radiological Dating Laboratory
The Norwegian Institute of Technology
Trondheim, Norway

Reidar Nydal was born in 1925 in Flora, Sogn og Fjordane, on the west coast of Norway. Reidar studied physics and mathematics at the University of Oslo and was awarded a master's degree in theoretical physics in 1953. That same year Reidar began work at the newly formed Radiological Dating Laboratory at the Norwegian Institute of Technology (NTH) in Trondheim under the direction of Professor Sverre Westin. In 1960, Reidar became head of the Radiological Dating Laboratory and he continued to direct the laboratory until one year before his retirement in 1995. Reidar received a doctorate from the University of Oslo in 1968, having written a dissertation entitled "An Investigation of Radiocarbon from Nuclear Tests".

Since 1986, Reidar has also served as a professor in the Physics Department at the NTH, where his research has focused on radiological dating. In addition, Reidar's research has focused on the global carbon cycle and the distribution of ^{14}C from nuclear tests in the atmosphere. To facilitate their work, Reidar and other staff (most notably Knut Lövseth) established a worldwide sampling program in the early 1960s to collect measurements in the atmosphere and ocean. Some of these stations have continued collecting measurements for approximately 30 years. Such extended time series are critical for testing global climate models based on CO_2 . During his distinguished career, Reidar has published 56 radiocarbon-related manuscripts, with roughly half of them related to the carbon cycle and climate. In November 1995, Reidar received the King Harald V gold medal for his radiocarbon work.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

CARBON-14 MEASUREMENTS IN ATMOSPHERIC CO₂ FROM NORTHERN AND SOUTHERN HEMISPHERE SITES, 1962-1993

Contributed by

Reidar Nydal and Knut Lövseth
Radiological Dating Laboratory
The Norwegian Institute of Technology
Trondheim, Norway

9845850

Prepared by

Virgene Zumbrunn* and Thomas A. Boden
Carbon Dioxide Information Analysis Center
World Data Center-A for Atmospheric Trace Gases
Oak Ridge National Laboratory
Oak Ridge, Tennessee

*ORISE summer intern

953 2500

Environmental Sciences Division
Publication No. 4582

Date Published: November 1996

Prepared for the
Global Change Research Program
Environmental Sciences Division
Office of Health and Environmental Research
U.S. Department of Energy
Budget Activity Number KP 1204010

MASTER

Prepared by the
Carbon Dioxide Information Analysis Center
World Data Center-A for Atmospheric Trace Gases
OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37831-6335
managed by
LOCKHEED MARTIN ENERGY RESEARCH CORP.
for the
U.S. DEPARTMENT OF ENERGY
under contract DE-AC05-96OR22464

SET 2AM

CONTENTS

| | <u>Page</u> |
|--|-------------|
| LIST OF FIGURES | v |
| LIST OF TABLES | vii |
| ABSTRACT | ix |
| PART 1: OVERVIEW | 1 |
| 1. NAME OF THE NUMERIC DATA PACKAGE | 3 |
| 2. PRINCIPAL INVESTIGATORS | 3 |
| 3. KEYWORDS | 3 |
| 4. BACKGROUND INFORMATION | 3 |
| 5. SAMPLING AND CALCULATION | 6 |
| 6. APPLICATIONS OF THE DATA | 7 |
| 7. DATA LIMITATIONS AND RESTRICTIONS | 7 |
| 8. REFERENCES | 9 |
| 9. DATA-PROCESSING ACTIVITIES AND QUALITY ASSURANCE CHECKS PERFORMED BY CDIAC | 10 |
| 10. HOW TO OBTAIN THE DATA AND DOCUMENTATION | 11 |
| PART 2: CONTENT AND FORMAT OF DATA FILES | 13 |
| 11. LISTING OF FILES PROVIDED | 14 |
| 12. FILE DESCRIPTIONS | 16 |
| 13. LISTING OF THE FORTRAN 77 DATA RETRIEVAL PROGRAMS | 21 |
| 14. LISTING OF THE SAS® DATA RETRIEVAL PROGRAMS | 23 |
| 15. PARTIAL LISTINGS OF DATA FILES | 24 |

| | |
|--|----|
| 16. VERIFICATION OF DATA TRANSPORT | 26 |
|--|----|

| | |
|---|-----|
| APPENDIX A: FIGURES SHOWING TROPOSPHERIC AND STRATOSPHERIC ¹⁴ C MEASUREMENTS IN ATMOSPHERIC CO ₂ FROM 14 NORTHERN AND SOUTHERN HEMISPHERE SITES | A-1 |
|---|-----|

| | |
|---|-----|
| APPENDIX B: REPRINT OF PERTINENT LITERATURE | B-1 |
|---|-----|

Tracing Bomb ¹⁴C in the Atmosphere 1962-1980, by R. Nydal and K. Lövseth. 1983.

LIST OF FIGURES

| <u>Figure</u> | <u>Page</u> |
|--|-------------|
| 1. Sampling network for tropospheric ^{14}C measurements in atmospheric CO_2 | 4 |
| 2. Corrected ^{14}C measurements from air samples collected at five Norwegian sites from 1963 to 1970 | 8 |
| A-1. Corrected ^{14}C measurements from air samples collected at Kapp Linné, Fruholmen, Gråkallen, and Vassfjellet | A-3 |
| A-2. Corrected ^{14}C measurements from air samples collected at Lindesnes, Norway; Santiago de Compostela, Spain; and Rehovot, Israel | A-4 |
| A-3. Corrected ^{14}C measurements from air samples collected at Izaña, Canary Islands; Mas Palomas, Grand Canary Island; Dakar, Senegal; Dakar-Fann, Senegal; and N'Djamena, Chad | A-5 |
| A-4. Corrected ^{14}C measurements from air samples collected at Debre Zeit, Ethiopia; Fianarantsoa, Madagascar; and by aircraft flying in the low stratosphere above Trondheim, Norway | A-6 |

LIST OF TABLES

| <u>Table</u> | <u>Page</u> |
|--|-------------|
| 1. Summary of the tropospheric $^{14}\text{CO}_2$ sampling sites | 5 |
| 2. Characteristics of numeric variables in the file containing carbon isotope data from all 14 sampling locations | 26 |
| 3. Characteristics of numeric variables in the file containing stratospheric carbon isotope data derived from air samples collected by aircraft during 1965 | 26 |

ABSTRACT

Nydal, R., and K. Lövseth. 1996. *Carbon-14 Measurements in Atmospheric CO₂ from Northern and Southern Hemisphere Sites, 1962–1993*. ORNL/CDIAC-93, NDP-057. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee. 67 pp. doi: 10.3334/CDIAC/atg.ndp057

In the 1960s, thermonuclear bomb tests released significant pulses of radioactive ^{14}C into the atmosphere. This major perturbation allowed scientists to study the dynamics of the global carbon cycle by measuring and observing rates of isotopic exchange. The Radiological Dating Laboratory at the Norwegian Institute of Technology performed ^{14}C measurements in atmospheric CO₂ from 1962 to 1993 at a network of ground stations in the Northern and Southern hemispheres. These measurements were supplemented during 1965 with high-altitude (9–12.6 km) air samples collected using aircraft from the Norwegian Air Force. The resulting database, coupled with other ^{14}C data sets, provides a greater understanding of the dynamic carbon reservoir and a crude picture of anomalous sources and sinks at different geographical latitudes. This database is outstanding for its inclusion of early ^{14}C measurements, broad spatial coverage of sampling, consistency of sampling method, and $\Delta^{14}\text{C}$ calculation results corrected for isotopic fractionation and radioactive decay. This database replaces previous versions published by the authors and the Radiological Dating Laboratory.

Fourteen stations spanning latitudes from Spitsbergen (78° N) to Madagascar (21° S) were used for sampling during the lifetime of the Norwegian program. Some of the stations have data for only a brief period, while others have measurements through 1993. Sampling stations subject to local industrial CO₂ contamination were avoided. The sites have sufficient separation to describe the latitudinal distribution of ^{14}C in atmospheric models. The sampling procedure for all the surface (10–2400 m asl) ^{14}C measurements in this database consisted of quantitative absorption of atmospheric CO₂ in carbonate-free 0.5 N NaOH solution. The ^{14}C measurements were made in a CO₂ proportional counter and calculated ($\delta^{14}\text{C}$) as per mil excess above the normal ^{14}C level defined by the US National Institute of Standards and Technology (NIST). Atmospheric ^{14}C content is finally expressed as $\Delta^{14}\text{C}$, which is the relative deviation of the measured ^{14}C activity from the NIST oxalic acid standard activity, after correction for isotopic fractionation and radioactive decay related to age. The data are organized by sampling station, and each record of the database contains the sampling dates; values for ^{14}C excess ($\delta^{14}\text{C}$) relative to the NIST standard, fractionation ^{13}C ($\delta^{13}\text{C}$) relative to the Pee Dee Belemnite (PDB) standard, and corrected ^{14}C ($\Delta^{14}\text{C}$) excess; and the standard deviation for $\Delta^{14}\text{C}$. The $\Delta^{14}\text{C}$ calculation results presented here are thus corrected for isotopic fractionation and radioactive decay, and constitute the final product of a research effort that has spanned three decades.

The $\Delta^{14}\text{C}$ station data show a sharp increase in tropospheric radiocarbon levels in the early 1960s and then a decline after the majority of nuclear tests came to an end on August 5, 1963 (Test Ban Treaty). The sharp peaks in tropospheric radiocarbon in the early 1960s are more pronounced in the Northern Hemisphere, reflecting the location of most atomic weapons tests. The measurements show large seasonal variations in the $\Delta^{14}\text{C}$ level during the early 1960s mainly as a result of springtime transport of bomb ^{14}C from the stratosphere. During the 1970s, the seasonal variations are smaller and due partly to seasonal variations in CO₂ from fossil-fuel emissions. The rate of decrease of atmospheric radiocarbon provides a check on the exchange constants of the atmosphere and ocean.

This report and all data it describes are available from the Carbon Dioxide Information Analysis Center (CDIAC) without charge. The Nydal and Lövseth atmospheric ^{14}C database comprises 21 data files totaling 0.2 megabytes in size. The following report describes the sampling methods and analysis. In addition, the report includes a complete discussion of CDIAC's data-processing efforts, the contents and format of the data files, and a reprint of a Nydal and Lövseth journal article.

PART 1
OVERVIEW

1. NAME OF THE NUMERIC DATA PACKAGE

Carbon-14 Measurements in Atmospheric CO₂ from Northern and Southern Hemisphere Sites,
1962–1993

2. PRINCIPAL INVESTIGATORS

Reidar Nydal and Knut Lövseth
Radiological Dating Laboratory
The Norwegian Institute of Technology
Trondheim, Norway

3. KEYWORDS

Radiocarbon, ¹⁴C, atmospheric CO₂, quantitative NaOH absorption, carbon isotopes, bomb ¹⁴C, Δ¹⁴C, δ¹⁴C, δ¹³C, troposphere, stratosphere, aircraft sampling

4. BACKGROUND INFORMATION

Atmospheric carbon dioxide (CO₂) contains, besides ¹²C (~99%), small amounts of the carbon isotopes ¹³C and ¹⁴C. Only ¹⁴C is radioactive with a half-life of 5730 years. In the atmosphere, ¹⁴C occurs principally as ¹⁴CO₂ and is usually produced by nuclear reactions between cosmic ray neutrons and the nitrogen atoms of the air (Libby 1952). Solar (heliomagnetic), geomagnetic, and ocean forcing all play a role in atmospheric ¹⁴CO₂ (Stuiver and Braziunas 1993). Before nuclear bomb testing during the 1950s and early 1960s, radiocarbon levels in the atmosphere had been in decline. Suess (1955) demonstrated that forests grown between 1930 and 1950 had Δ¹⁴C values 20–40 per mil below those of pre-1890 woods as a result of anthropogenic ¹⁴C-free CO₂ emissions since the Industrial Revolution. Nuclear bomb testing has since swamped the "Suess Effect".¹ After a series of atomic weapons tests were conducted at higher northern latitudes, notably Novaya Zemlya in the former Soviet Union, scientists at the Radiological Dating Laboratory in Norway seized the opportunity to conduct CO₂ sampling to trace the distribution and flux of bomb ¹⁴C with funding support from the Norwegian Research Council. In the early 1960s, several ground sampling sites were established in Norway. These Norwegian study sites gradually expanded to a global network of 14 stations spanning latitudes from Spitsbergen (78° N) to Madagascar (21° S) (Fig. 1; Table 1). Sites possibly contaminated by local CO₂ sources were avoided. Additional sites were chosen to improve spatial coverage so that the

¹The relative change in the ¹⁴C or ¹³C ratio of any carbon pool or reservoir caused by the addition of fossil-fuel CO₂ to the atmosphere. Fossil fuels are devoid of ¹⁴C because of the radioactive decay of ¹⁴C to ¹⁴N during long underground storage and are depleted in ¹³C because of isotopic fractionation long ago during photosynthesis by the plants that were the precursors of the fossil fuels. Carbon dioxide produced by the combustion of fossil fuels is thus virtually free of ¹⁴C and depleted in ¹³C. The term "Suess Effect" originally referred to the dilution of the ¹⁴C/C ratio in atmospheric CO₂ but the definition has been extended to both the ¹⁴C and ¹³C ratios in any pool or reservoir of the carbon cycle resulting from human disturbances (Keeling 1979).




Figure 1. Sampling network for tropospheric ^{14}C measurements in atmospheric CO_2

Table 1. Summary of the tropospheric $^{14}\text{CO}_2$ sampling sites

| Site code | Site | Country | Agency | Latitude | Longitude | Elevation | Period of record |
|-----------|------------------------|------------|-----------------------------------|-----------|-----------|-----------|------------------|
| V | Kapp Linné | Norway | Telegrafstyret | 78° 04' N | 13° 38' E | 10 m | 1963-64 |
| F | Fruholmen | Norway | Fruholmen lighthouse | 71° 06' N | 23° 59' E | 70 m | 1962-93 |
| K | Gråkallen | Norway | Norwegian Institute of Technology | 63° 25' N | 10° 15' E | 450 m | 1962 |
| Kv | Vassfjellet | Norway | Norwegian Institute of Technology | 63° 16' N | 10° 22' E | 700 m | 1962-63 |
| L | Lindesnes | Norway | Lindesnes lighthouse | 57° 59' N | 07° 04' E | 50 m | 1963-64 |
| S | Santiago de Compostela | Spain | Local airport | 42° 53' N | 08° 26' W | 360 m | 1963-66 |
| IZ | Izaña | Spain | Observatorio Astronomica, Izaña | 28° 22' N | 16° 30' W | 2400 m | 1963-67, 1976-90 |
| M | Mas Palomas | Spain | NASA | 27° 45' N | 15° 40' W | 10-100 m | 1963-73 |
| R | Rehovot | Israel | Weismann Institute of Science | 31° 50' N | 34° 50' E | 100 m | 1967-68 |
| D | Dakar | Senegal | University of Dakar-Fann | 14° 33' N | 17° 07' W | 10 m | 1963-64 |
| DF | Dakar-Fann | Senegal | University of Dakar-Fann | 14° 41' N | 17° 28' W | 25 m | 1964-68 |
| TC | N'Djamena | Chad | ORSTOM de Fort-Lamy | 12° 08' N | 14° 57' E | 300 m | 1966-76 |
| E | Debre Zeit | Ethiopia | Ethiopian Evangelical College | 8° 40' N | 38° 58' E | 1900 m | 1963-69 |
| G | Fianarantsoa | Madagascar | Missionary station | 21° 27' S | 47° 05' E | 1100 m | 1964-78 |

distribution of radiocarbon could be characterized with respect to latitude. During 1965 the ground-level measurements were supplemented with measurements made on air samples collected from the lower stratosphere (9000–12600 m) using aircraft from the Norwegian Air Force. This data document provides the surface ^{14}C measurements made by the Radiological Dating Laboratory at this global sampling network and the high-altitude measurements made during 1965. The $\Delta^{14}\text{C}$ data presented here are corrected for isotopic fractionation and radioactive decay, and constitute the final product of a research effort that has spanned three decades. The earliest measurements date back to 1962, and a few sites have data through 1993. Earlier versions of this database have been presented by Nydal (1966, 1968, 1993), Nydal and Lövseth (1983), and Nydal et al. (1971).

5. SAMPLING AND CALCULATION

All ground-level air samples represented in this database were collected by dynamic quantitative absorption of atmospheric CO_2 in carbonate-free 0.5 *N* sodium hydroxide (NaOH) solution. During the collection process, a dish of NaOH was exposed to the air for 4 to 7 days. Following exposure, the samples were treated with hydrochloric acid to regenerate the carbon dioxide. After a purification procedure, the CO_2 was analyzed in a proportional counter to assess ^{14}C content. Typically, a counter has gas volumes of 1–2 L and operates at 2 atm pressure. Until 1981 a counting time of 2 days (48 hours) was often used (Nydal and Lövseth 1983), but a counting time of 4 days was used later in order to obtain higher precision.

High-altitude samples were collected in the Trondheim area during 1965 using aircraft from the Norwegian Air Force. Flights originated from Örlandet Airport and covered a region bounded by 62–65° N and 6–10° E, at altitudes of 9000–12600 m. Samples were collected using a filter with a 1.4 kg molecular sieve that absorbs CO_2 from the penetrating air. The filter container was placed under the wing of the plane and opened and closed with electrically-operated shutters at both ends. With the plane traveling at a speed of approximately 900 km/hr the filter shutters were opened 15 minutes to allow sufficient amounts of CO_2 to be absorbed.

After determination of the ^{14}C content in a proportional counter, the $\delta^{14}\text{C}$ values were calculated as per mil excess above the normal ^{14}C level defined by the National Institute of Standards and Technology (NIST, formerly the US National Bureau of Standards). Atmospheric ^{14}C content is finally expressed as $\Delta^{14}\text{C}$, which is the relative deviation of the measured ^{14}C activity from the NIST oxalic acid standard activity, after correction for isotopic mass fractionation and radioactive decay related to age (Stuiver and Polach 1977). $\Delta^{14}\text{C}$ is expressed in per mil (not as a percentage) and calculated using the following two steps:

$$\Delta^{14} = \delta^{14}\text{C} - 2(\delta^{13}\text{C} + 25)(1 + \delta^{14}\text{C}/1000) \quad (1)$$

$$\Delta^{14}\text{C} = \Delta^{14} + 1000(e^{\lambda(1950-t)} - 1) \quad (2)$$

In step 1, the ^{14}C excess (Δ^{14}) is only corrected for isotopic mass fractionation. $\delta^{14}\text{C}$ represents the uncorrected (for decay) ^{14}C relative to the NIST standard and $\delta^{13}\text{C}$ represents the deviation in the $^{13}\text{C}/^{12}\text{C}$ ratio relative to the Pee Dee Belemnite (Pee Dee River, South Carolina) standard. The $^{13}\text{C}/^{12}\text{C}$ ratio was measured by mass spectrometry. By themselves, the $\delta^{13}\text{C}$ values represent data that are highly influenced by fractionation in NaOH in the absorbing dish. In the earlier presentation of the ^{14}C data from the Radiological Dating Laboratory only step 1 was used, and the decay of the NIST ^{14}C reference standard after 1950 was not taken into account (Nydal and Lövseth 1983). Applying the

approximate formula in step 2, where λ is $1/8267$ years ($T/\ln 2$ where T is the 5730 year half-life of ^{14}C) and t is the year of sampling, this small decay (2–5 per mil) has now been applied to all the $\Delta^{14}\text{C}$ data appearing in this database.

For a further discussion of the sampling methods and each sampling site, please see the reprint of Nydal and Lövseth (1983) in Appendix B. For additional information on the high-altitude sampling, see Hagemann et al. (1965), Nydal (1966), and Nydal and Lövseth (1983). For additional information on the sampling sites, radiocarbon standards, and carbon isotope measurements, see Nydal (1966, 1968), Craig (1961), and Levin et al. (1980).

6. APPLICATIONS OF THE DATA

These measurements were originally made to assess potential human health risks from radioactive fallout in the atmosphere. Fortunately it was also recognized that the radioactive isotopes already injected into the atmosphere could be useful tracers in geophysical research. These ^{14}C measurements document the time lag of isotopic sea-air exchange and chronicle a long-term decline in $\Delta^{14}\text{CO}_2$ at Northern and Southern hemisphere sites. These measurements complement other radiocarbon data sets (Broecker and Olson 1959; Levin et al. 1985, 1995; Manning et al. 1990; Olsson 1993) and support the pronounced twentieth-century $\Delta^{14}\text{C}$ reduction reported by Stuiver and Quay (1981) due to ^{14}C -free carbon dioxide released by fossil-fuel combustion.

This data set is unique for its inclusion of measurements in the early 1960s that trace the bomb ^{14}C released by a series of nuclear tests. All measurements at the global network of sites were taken by dynamic quantitative absorption of CO_2 in NaOH solution. This consistency in sampling technique over a period of three decades is valuable to researchers and modelers.

The $\Delta^{14}\text{C}$ tropospheric data show a sharp increase in tropospheric radiocarbon levels in the early 1960s and then a decline after the majority of nuclear tests came to an end on August 5, 1963 through the Test Ban Treaty. The sharp peaks in tropospheric radiocarbon in the early 1960s are more pronounced in the Northern Hemisphere, reflecting the location of most atomic weapons tests. The measurements show large seasonal variations in the $\Delta^{14}\text{C}$ level in the early 1960s, mainly as a result of springtime transport of bomb ^{14}C from the stratosphere (Fig. 2). During the 1970s, the seasonal variations are smaller and are partly due to seasonal variations in CO_2 from fossil-fuel emissions. The rate of decrease of atmospheric radiocarbon provides a check on the exchange constants of the atmosphere and ocean.

7. DATA LIMITATIONS AND RESTRICTIONS

Please note that the $\delta^{13}\text{C}$ values provided in this database are not representative of atmospheric values. In this database the ^{13}C data serve as guides to the ^{14}C measurements and were used to calculate corrected ^{14}C . It should also be noted that the $^{13}\text{C}/^{12}\text{C}$ ratio was not measured for every sample. In these instances where the $^{13}\text{C}/^{12}\text{C}$ ratio was not measured, a mean value is given for $\delta^{13}\text{C}$ and identified by a flag code. These mean values were calculated from 20–40 ground-level measurements or 4 high-altitude measurements.




Figure 2. Corrected ^{14}C measurements from air samples collected at five Norwegian sites from 1963 to 1970

8. REFERENCES

- Broecker, W. S., and E. A. Olson. 1959. Lamont Radiocarbon Measurements VI*. *American Journal of Science Radiocarbon Supplement* 1:111–32.
- Craig, H. 1961. Mass-spectrometer analyses of radiocarbon standards. *Radiocarbon* 3:1–3.
- Hagemann, F.T., J. Gray, Jr., and L. Machta. 1965. Carbon 14 measurements in the atmosphere - 1953 to 1964. U.S. Atomic Energy Commission Report. HASL-159, 124 pgs.
- Keeling, C.D. 1979. The Suess Effect: ^{13}C - ^{14}C Interrelations. *Environment International* 2:229–300.
- Levin, I., K. O. Münnich, and W. Weiss. 1980. The effect of anthropogenic CO_2 and ^{14}C sources on the distribution of ^{14}C in the atmosphere. *Radiocarbon* 22:379–91.
- Levin, I., B. Kromer, H. Schoch-Fischer, M. Bruns, M. Münnich, D. Berdau, J. C. Vogel, and K. O. Münnich. 1985. 25 years of tropospheric ^{14}C observations in central Europe. *Radiocarbon* 27(1):1–19.
- Levin, I., R. Graul, and N. B. A. Trivett. 1995. Long-term observations of atmospheric CO_2 and carbon isotopes at continental sites in Germany. *Tellus* 47(B):23–34.
- Libby, W.F. 1952. Radiocarbon dating. The University of Chicago Press, Chicago, Illinois, USA 161 pgs.
- Manning, M. R., D. C. Lowe, W. H. Melhuish, R. J. Sparks, G. Wallace, C. A. M. Brenninkmeijer, and R. C. McGill. 1990. The use of radiocarbon measurements in atmospheric studies. *Radiocarbon* 32(1):37–58.
- Nydal, R. 1966. Variation in C^{14} concentration in the atmosphere during the last several years. *Tellus* 18:271–79.
- Nydal, R. 1968. Further investigation on the transfer of radiocarbon in nature. *Journal of Geophysical Research* 73:3617–35.
- Nydal, R. 1993. Application of bomb ^{14}C as a tracer in the global carbon cycle. *Trends in Geophysical Research* 2:355–64.
- Nydal, R., and K. Lövseth. 1983. Tracing bomb ^{14}C in the atmosphere, 1962–1980. *Journal of Geophysical Research* 88:3621–42.
- Nydal, R., K. Lövseth, and O. Syrstad. 1971. Bomb ^{14}C in the human population. *Nature* 232:418–21.

- Olsson, I. U. 1993. A ten-year record of different levels of the ^{14}C activities over Sweden and the Arctic. *Tellus* 45(B):479–81.
- Stuiver, M., and H. A. Polach. 1977. Discussion: Reporting of ^{14}C data. *Radiocarbon* 19:355–63.
- Stuiver, M., and P. D. Quay. 1981. Atmospheric ^{14}C changes resulting from fossil fuel CO_2 release and cosmic ray flux variability. *Earth and Planetary Science Letters* 53:349–62.
- Stuiver, M., and T. F. Braziunas. 1993. Sun, ocean, climate and atmospheric $^{14}\text{CO}_2$: An evaluation of causal and spectral relationships. *The Holocene* 3(4):289–305.
- Suess, H. E. 1955. Radiocarbon concentration in modern wood. *Science* 122:415–17.

9. DATA-PROCESSING ACTIVITIES AND QUALITY ASSURANCE CHECKS PERFORMED BY CDIAC

CDIAC is committed to the quality assurance (QA) of data before distribution. In order to provide scientists and researchers with high-quality data, CDIAC reviews the data it receives for consistent formatting, completeness, reasonableness, and accuracy. Each review involves programming that is specific to the needs of each data set. These efforts are indicative of CDIAC's mission to provide the international scientific community with high-quality, well-documented data.

DATA-PROCESSING

CDIAC obtained the Nydal and Lövseth database as an electronic-mail message from Reidar Nydal. CDIAC staff edited the message and created the ASCII data files for each site. Working copies of the files were created and processed in the following ways:

1. The original ASCII files were reformatted into a consistent format and combined in a single file with data from all sites.
2. Both SAS® and FORTRAN 77 codes were written and are available to the user for reading any of the ASCII data files.

QA CHECKS

1. Each file was checked to ensure formatting consistency and to confirm the absence of missing data entries.
2. Checks were performed to confirm the uniqueness of each laboratory reference code and to ensure that sampling intervals were formatted consistently.
3. Mean, minimum, and maximum values for $\delta^{14}\text{C}$, $\delta^{13}\text{C}$, and corrected $\Delta^{14}\text{C}$ data at each site were calculated and assessed for reasonableness.

SAS® is the registered trademark of SAS Institute, Inc., Cary, North Carolina, 27511, USA.

4. All $\delta^{14}\text{C}$, $\delta^{13}\text{C}$, and corrected $\Delta^{14}\text{C}$ measurements were plotted and assessed for reasonableness.
5. Any inconsistencies or suspect measurements were reported to and addressed by the investigators before the data set was released.

10. HOW TO OBTAIN THE DATA AND DOCUMENTATION

The Nydal and Lövseth database is available in machine-readable form from CDIAC without charge. The database may also be downloaded from CDIAC's anonymous File Transfer Protocol (FTP) site, by using the address, login instructions, and FTP commands listed below. In addition, the database is available on 9-track magnetic tape, 8-mm tape, or IBM- or Macintosh-formatted floppy diskettes. For tape requests, please include preferred tape specifications (i.e., 1600 or 6250 BPI for 9-track tapes and 8200 or 8500 format for 8-mm tapes, labeled or nonlabeled, ASCII or EBCDIC characters, variable or fixed-record lengths). Requests without tape specifications will be filled on 9-track, 6250 BPI, nonlabeled tapes with file attributes shown in Sect. 11.

This documentation is available only from CDIAC. Electronic versions of the documentation are available via the World Wide Web from CDIAC's home page (<http://cdiac.esd.ornl.gov>). Requests for printed copies of the documentation should be addressed to

Carbon Dioxide Information Analysis Center
World Data Center-A for Atmospheric Trace Gases
Oak Ridge National Laboratory
Post Office Box 2008
Oak Ridge, TN 37831-6335, USA

The tapes, diskettes, and documentation may also be ordered by telephone, facsimile, or electronic mail:

Telephone: (423) 574-3645 or (423) 574-0390

Fax: (423) 574-2232

Electronic mail: cdiac@ornl.gov

FTP access: [ftp cdia.esd.ornl.gov](ftp://cdiac.esd.ornl.gov) (or 128.219.24.36)

Enter **anonymous** at the userid prompt

Enter your e-mail address as the password

Change the directory to **/pub/ndp057** (i.e., `ftp> cd /pub/ndp057`)

Retrieve all files (i.e., `ftp> mget *`)

NOTE: When using these radiocarbon data in a presentation or publication, **PLEASE** acknowledge the principal investigators, Reidar Nydal and Knut Lövseth, and the Norwegian Institute of Technology!

PART 2

CONTENT AND FORMAT OF DATA FILES

11. LISTING OF FILES PROVIDED

The following is a list of the files that compose the Nydal and Lövseth atmospheric carbon isotope database and that are distributed by CDIAC along with this documentation. These files are available on a variety of media (see Sect. 10, "How To Obtain the Data and Documentation"). This listing has been tailored to reflect a 9-track magnetic tape request. The record formats, block sizes, and record lengths shown are the defaults for those not specifying these parameters when requesting machine-readable data files on 9-track magnetic tape. The default tape density, labeling, and characters are 6250 BPI, nonlabeled, and ASCII, respectively.

| File number, description, and name (in parentheses) | Logical records | FTP file size (kB) | Record format ^a | Block size | Record length |
|--|--------------------|-----------------------|-------------------------------|---------------|------------------|
| 1. General descriptive information file (ndp057.doc) | 255 | 13.3 | FB | 8000 | 100 |
| 2. FORTRAN 77 data retrieval code to read and print the files containing the atmospheric carbon isotope records (files 4–18) (ndp057.for) | 21 | 0.7 | FB | 8000 | 80 |
| 3. SAS [®] data retrieval code to read and print files 4–18 (ndp057.sas) | 12 | 0.4 | FB | 8000 | 80 |
| Data files containing $\delta^{14}\text{C}$, $\delta^{13}\text{C}$, and $\Delta^{14}\text{C}$ estimates in atmospheric CO_2 from the following Northern and Southern Hemisphere sites | | | | | |
| 4. Kapp Linné, Spitsbergen, Norway (V.asc) | 59 | 1.6 | FB | 6100 | 61 |
| 5. Fruholmen, Nordkapp, Norway (F.asc) | 531 | 32.5 | FB | 6100 | 61 |
| 6. Gråkallen, Trondheim, Norway (K.asc) | 44 | 2.6 | FB | 6100 | 61 |
| 7. Vassfjellet, Trondheim, Norway (KV.asc) | 32 | 1.8 | FB | 6100 | 61 |
| 8. Lindesnes, Norway (L.asc) | 64 | 3.8 | FB | 6100 | 61 |
| 9. Santiago de Compostela, Spain (S.asc) | 104 | 6.3 | FB | 6100 | 61 |
| 10. Izaña, Tenerife, Canary Islands (IZ.asc) | 308 | 19.3 | FB | 6100 | 61 |
| 11. Mas Palomas, Grand Canary Island (M.asc) | 183 | 11.4 | FB | 6100 | 61 |
| 12. Rehovot, Israel (R.asc) | 28 | 1.6 | FB | 6100 | 61 |

| File number, description, and name (in parentheses) | Logical records | FTP file size (kB) | Record format ^a | Block size | Record length |
|---|--------------------|-----------------------|-------------------------------|---------------|------------------|
| 13. Popenguine, Dakar, Senegal (D.asc) | 48 | 2.8 | FB | 6100 | 61 |
| 14. Dakar-Fann, Senegal (DF.asc) | 84 | 5.1 | FB | 6100 | 61 |
| 15. N'Djamena, Chad (TC.asc) | 119 | 7.2 | FB | 6100 | 61 |
| 16. Debre Zeit, Ethiopia (E.asc) | 84 | 5.2 | FB | 6100 | 61 |
| 17. Fianarantsoa, Madagascar (G.asc) | 185 | 11.4 | FB | 6100 | 61 |
| 18. Single data file containing the $\delta^{14}\text{C}$, $\delta^{13}\text{C}$, and $\Delta^{14}\text{C}$ estimates in atmospheric CO_2 from all 14 sites (nydal.asc) | 1635 | 101.8 | FB | 6100 | 61 |
| 19. FORTRAN 77 data retrieval code to read and print the file containing the stratospheric $\delta^{14}\text{C}$, $\delta^{13}\text{C}$, and $\Delta^{14}\text{C}$ estimates in atmospheric CO_2 derived from air samples collected by aircraft during 1965 from the Trondheim area (62–65° N, 6–10° E) (strat.for) | 26 | 0.9 | FB | 8000 | 80 |
| 20. SAS [®] data retrieval code to read and print file 21 (strat.sas) | 13 | 0.5 | FB | 8000 | 80 |
| 21. Stratospheric $\delta^{14}\text{C}$, $\delta^{13}\text{C}$, and $\Delta^{14}\text{C}$ estimates in atmospheric CO_2 derived from air samples collected by aircraft during 1965 from the Trondheim area (62–65° N, 6–10° E) (strat.dat) | 27 | 1.9 | FB | 7600 | 76 |
| TOTAL | 3862 | 232.1 | | | |

^aFixed-block record format.

[®]SAS is the registered trademark of SAS Institute, Inc., Cary, North Carolina, 27511, USA.

12. FILE DESCRIPTIONS

This section describes the content and format of each of the 21 files that compose this numeric data package (NDP) and the Nydal and Lövseth atmospheric carbon isotope database.

DATA FILES

This NDP consists of the following 21 files:

- a descriptive file that contains an overview of the Nydal and Lövseth atmospheric carbon isotope database and detailed descriptions of the data files (**ndp057.doc**);
- one FORTRAN 77 data retrieval routine to read and print any of the 15 data files containing estimates of $\delta^{14}\text{C}$, $\delta^{13}\text{C}$, and $\Delta^{14}\text{C}$ in atmospheric CO_2 from air samples collected at 14 sites (**ndp057.for**);
- one SAS[®] data retrieval routine to read and print any of the 15 data files (**ndp057.sas**);
- one ASCII data file containing estimates of $\delta^{14}\text{C}$, $\delta^{13}\text{C}$, and $\Delta^{14}\text{C}$ in atmospheric CO_2 from all 14 sites (**nydal.asc**);
- 14 ASCII data files (one for each sampling location) containing estimates of $\delta^{14}\text{C}$, $\delta^{13}\text{C}$, and $\Delta^{14}\text{C}$ in atmospheric CO_2 (*.asc — see description of "LABREF" on page 15),
- one FORTRAN 77 data retrieval routine to read and print the file containing estimates of $\delta^{14}\text{C}$, $\delta^{13}\text{C}$, and $\Delta^{14}\text{C}$ in atmospheric CO_2 collected in the lower stratosphere from aircraft during 1965 (**strat.for**);
- one SAS[®] data retrieval routine (**strat.sas**) to read and print **strat.dat**; and
- one ASCII data file containing estimates of $\delta^{14}\text{C}$, $\delta^{13}\text{C}$, and $\Delta^{14}\text{C}$ in atmospheric CO_2 collected in the lower stratosphere from aircraft during 1965 (**strat.dat**)

DATA FILE FORMATS (*.ASC)

This NDP contains 15 data files that provide 1619 estimates of $\delta^{14}\text{C}$, $\delta^{13}\text{C}$, and $\Delta^{14}\text{C}$ in atmospheric CO_2 from 14 locations. The period of record differs by station. The earliest measurements were made in 1962, and the latest estimates are from 1993. Most stations have records only for the 1960s. All of the data files have the same format and none contain missing values. Some $\delta^{13}\text{C}$ values are flagged as mean values; users are urged to pay attention to these flags before analyzing or interpreting the corresponding values.

Each file is sorted by sample collection date and may be read with the following FORTRAN 77 code:

```
character labref*9, samdat*13, flag13*1
real delC14, delC13, corC14
open(unit=5, file='F.asc')
open(unit=6)
read(5,10)
10  format(//////////)
13  continue
    read (5,15, end=99) labref, week, samdat, delC14, delC13, flag13,
1  corC14, sigma
15  format (A9,1X,I4,2X,A13,4X,F6.1,2X,F5.1,A1,2X,F6.1,4X,I2)
```

The following SAS® input statement may also be used to read these files:

```
input labref $ 1-9 week 11-15 samdat $ 17-30 @34 delC14 7.1 @42 delC13 5.1
      flag13 $ 47 corC14 50-55 sigma 60-61;
```

Stated in tabular form, the contents include the following:

| Variable | Variable type | Variable format | Starting column | Ending column |
|----------|---------------|-----------------|-----------------|---------------|
| LABREF | Alphanumeric | 9 | 1 | 9 |
| WEEK | Numeric | 5 | 11 | 15 |
| SAMDAT | Alphanumeric | 14 | 17 | 30 |
| DELC14 | Numeric | 7.1 | 34 | 40 |
| DELC13 | Numeric | 5.1 | 42 | 46 |
| FLAG13 | Character | 1 | 47 | 47 |
| CORC14 | Numeric | 6.1 | 50 | 55 |
| SIGMA | Numeric | 2 | 60 | 61 |

where

LABREF is a seven-to-eight character site code and sample reference number (e.g., D_00034). The 5-digit sample reference number follows the site code and is separated from the site code by an underscore (i.e., "_") character. All laboratory reference numbers are unique. The site codes and the sampling stations they represent are as follows:

| | |
|----|---------------------------------------|
| V | Kapp Linné, Spitsbergen, Norway |
| F | Fruholmen, Nordkapp, Norway |
| K | Gråkallen, Trondheim, Norway |
| KV | Vassfjellet, Trondheim, Norway |
| L | Lindesnes, Norway |
| S | Santiago de Compostela, Spain |
| IZ | Izaña, Tenerife, Canary Islands |
| M | Mas Palomas, Grand Canary Island |
| R | Rehovot, Israel |
| D | Popenguine, Dakar, Senegal |
| DF | Université Dakar-Fann, Dakar, Senegal |
| TC | N'Djamena, Chad |
| E | Debre Zeit, Addis Ababa, Ethiopia |
| G | Fianarantsoa, Madagascar |

WEEK is the week the NaOH tray was exposed relative to the first week of 1963. Values range from -34 to 1591. Zero refers to the week of December 29, 1962 to

January 7, 1963. Negative values denote the number of weeks before 1963 (e.g., -34 refers to the week from April 30, 1962, to May 3, 1962).

- SAMDAT** is a thirteen-character field that represents the days, months, and years of the sampling period, typically 4–7 day periods. The beginning date (yy/mm/dd) and ending date (yy/mm/dd) are separated by a hyphen. The earliest sampling date is 620430-620503 (from Gråkallen) and the latest sampling period is 930621-930624 (from Fruholmen).
- DELC14** is the uncorrected (for decay) measurement of excess ^{14}C in atmospheric CO_2 expressed as $\delta^{14}\text{C}$ in per mil (‰) relative to the US National Institute of Standards and Technology (NIST, formerly the National Bureau of Standards) oxalic acid standard; 95% of the ^{14}C concentration in this standard represents the normal activity in recent wood for the year 1950 A.D. Values for DELC14 range from 108.5 to 1036.0.
- DELC13** represents the deviation in the $^{13}\text{C}/^{12}\text{C}$ ratio expressed in per mil (‰) relative to the Pee Dee Belemnite or (Pee Dee River, South Carolina) or PDB standard (Craig 1961). The $^{13}\text{C}/^{12}\text{C}$ ratio was measured by mass spectrometry. The $\delta^{13}\text{C}$ values provided are given as a guide for the calculation of the $\Delta^{14}\text{C}$ values. When fractionation was not measured, a mean value, normally based on 20–40 samples was used and is denoted by an asterisk (see description of "FLAG13"). Values for DELC13 range from -31.7 to -11.3.
- FLAG13** is a one-character flag code used to denote whether the DEL13C estimate was measured (blank) or represents a mean value (asterisk).
- CORC14** is the atmospheric ^{14}C content expressed as $\Delta^{14}\text{C}$ in per mil (not as a percentage). $\Delta^{14}\text{C}$ is corrected for isotopic fractionation through $\delta^{13}\text{C}$ (DELC13), and for radioactive decay of the ^{14}C reference standard (NIST). Values range from 110.4 to 1040.4.
- SIGMA** is the standard deviation for CORC14 expressed as an integer. Values range from 3 to 42.

DATA FILE FORMAT (STRAT.DAT)

This NDP contains one data file that provides eleven estimates of $\delta^{14}\text{C}$, $\delta^{13}\text{C}$, and $\Delta^{14}\text{C}$ in atmospheric CO_2 from the low stratosphere during 1965. High-altitude samples were collected in the Trondheim area using aircraft from the Norwegian Air Force. Flights originated from Örlandet Airport and covered the region bounded by 62–65° N and 6–10° E, at altitudes of 9000-12600 m. Samples were collected using a filter with a 1.4 kg molecular sieve that absorbs CO_2 from the penetrating air. The filter container was placed under the wing of the plane and opened and closed with electrically-operated shutters at both ends. With the plane traveling at a speed of approximately 900 km/hr the filter shutters were opened 15 minutes to allow sufficient amounts of CO_2 to be absorbed.

The file may be read with the following FORTRAN 77 code:

```

character sample*6, flag13*1
real delC13, corC14
open(unit=5,file='strat.dat')
open(unit=6)
read(5,10)
10 format(////////////////)
write(6,15)
15 format('Sample',3X,'Altitude',3X,'Date of Collection',6X,
1 'delta',6X,'delta',2X,'Corrected',4X,'Sigma',/,11X,'(m)',
2 6X,'(Year, Month, Day)',7X,'C14',8X,'C13',3X,'delta C14',
3 /)
20 continue
read(5,25, end=99) sample,alt,samdat,delC14,delC13,flag13,
1 corC14,sigma
25 format(A6,4X,I5,11X,I6,12X,I4,4X,F5.1,A1,4X,I4,8X,I2)

```

The following SAS® input statement may also be used to read this file:

```

input sample $ 1-6 alt 11-15 samdat 27-32 delC14 45-48 @53 delC13 5.1
flag13 $ 58 corC14 63-66 sigma 75-76;

```

Stated in tabular form, the contents include the following:

| Variable | Variable type | Variable format | Starting column | Ending column |
|----------|------------------|--------------------|--------------------|------------------|
| SAMPLE | Alphanumeric | 6 | 1 | 6 |
| ALT | Numeric | 5 | 11 | 15 |
| SAMDAT | Numeric | 6 | 27 | 32 |
| DELC14 | Numeric | 4 | 45 | 48 |
| DELC13 | Numeric | 5.1 | 53 | 57 |
| FLAG13 | Character | 1 | 58 | 58 |
| CORC14 | Numeric | 4 | 63 | 66 |
| SIGMA | Numeric | 2 | 75 | 76 |

where

SAMPLE is a six character site code and sample reference number (e.g., AS-001). The 3-digit sample reference number follows the site code and is separated from the site code by a hyphen. All laboratory reference numbers are unique. Laboratory reference numbers range from 001 to 013. The site code is always "AS" denoting aircraft sampling.

ALT is the altitude where the air sample was collected expressed in meters. Values range from 9000 to 12600.

| | |
|--------|---|
| SAMDAT | is a six-digit field that represents the day, month, and year of the sampling date expressed in a yymmdd format. Values range from 650407 to 651014. |
| DELC14 | is the uncorrected (for decay) measurement of excess ^{14}C in atmospheric CO_2 expressed as $\delta^{14}\text{C}$ in per mil (‰) relative to the US National Institute of Standards and Technology (NIST, formerly the National Bureau of Standards) oxalic acid standard; 95% of the ^{14}C concentration in this standard represents the normal activity in recent wood for the year 1950 A.D. Values for DELC14 range from 599 to 1718. |
| DELC13 | represents the deviation in the $^{13}\text{C}/^{12}\text{C}$ ratio expressed in per mil (‰) relative to the Pee Dee Belemnite (Pee Dee River, South Carolina) or PDB standard (Craig 1961). The $^{13}\text{C}/^{12}\text{C}$ ratio was measured by mass spectrometry. The $\delta^{13}\text{C}$ values provided are given as a guide for the calculation of the $\Delta^{14}\text{C}$ values. When fractionation was not measured, a mean value, based on 4 samples was used and is denoted by an asterisk (see description of "FLAG13"). Values for DELC13 range from -11.4 to -3.3. |
| FLAG13 | is a one-character flag code used to denote whether the DEL13C estimate was measured (blank) or represents a mean value (asterisk). Seven of the eleven DELC13 values are flagged with an asterisk. |
| CORC14 | is the atmospheric ^{14}C content expressed as $\Delta^{14}\text{C}$ in per mil (not as a percentage). $\Delta^{14}\text{C}$ is corrected for isotopic fractionation through $\delta^{13}\text{C}$ (DELC13), and for radioactive decay of the ^{14}C reference standard (NIST). Values range from 554 to 1626. |
| SIGMA | is the standard deviation for CORC14 expressed as an integer. Values range from 17 to 47. |

REFERENCES

- Craig, H. 1961. Mass-spectrometer analysis of radiocarbon standards. *American Journal of Science Radiocarbon Supplement* 3:1-3.
- Nydal, R., and K. Lövseth. 1983. Tracing bomb ^{14}C in the atmosphere 1962-1980. *Journal of Geophysical Research* 88:3621-42.

13. LISTING OF THE FORTRAN 77 DATA RETRIEVAL PROGRAMS

The following is a listing of the FORTRAN 77 data retrieval code (**ndp057.for**) written by the preparers to read and print any of the files containing the station ^{14}C data.

```
C This Fortran 77 program was written by VZ to read and write
C Nydal and Lovseth's C14 measurements
  character labref*9, samdat*13, flag13*1
  real delC14, delC13, corC14
  open(unit=5,file='F.asc')
  open(unit=6)
  read(5,10)
10  format(////////////////)
13  continue
  read (5,15, end=99) labref,week,samdat,delC14,delC13,flag13,
1  corC14,sigma
15  format(A9,1X,I4,2X,A13,4X,F6.1,2X,F5.1,A1,2X,F6.1,4X,I2)
  write(6,20) labref,week,samdat,delC14,delC13,flag13,
1  corC14,sigma
20  format(A9,1X,I4,2X,A13,4X,F6.1,2X,F5.1,A1,2X,F6.1,4X,I2)
  goto 13
99  continue
  close(unit=5)
  close(unit=6)
  stop
  end
```

The following is a listing of the FORTRAN 77 data retrieval code (**strat.for**) written by the preparers to read and print the file containing the stratospheric ^{14}C data.

```

C This Fortran 77 program was written by TAB to read and write
C Nydal and Lovseth's aircraft C14 measurements
  character sample*6, flag13*1
  real delC13, corC14
  open(unit=5,file='strat.dat')
  open(unit=6)
  read(5,10)
10  format(/////////)
  write(6,15)
15  format('Sample',3X,'Altitude',3X,'Date of Collection',6X,
1  'delta',6X,'delta',2X,'Corrected',4X,'Sigma',/,11X,'(m)',
2  6X,'(Year, Month, Day)',7X,'C14',8X,'C13',3X,'delta C14',
3  /)
20  continue
  read (5,25, end=99) sample,alt,samdat,delC14,delC13,flag13,
1  corC14,sigma
25  format(A6,4X,I5,11X,I6,12X,I4,4X,F5.1,A1,4X,I4,8X,I2)
  write(6,30) sample,alt,samdat,delC14,delC13,flag13,
1  corC14,sigma
30  format(A6,4X,I5,11X,I6,12X,I4,6X,F5.1,A1,4X,I4,8X,I2)
  goto 20
99  continue
  close(unit=5)
  close(unit=6)
  stop
  end

```

14. LISTING OF THE SAS® DATA RETRIEVAL PROGRAMS

The following is a listing of the SAS® data retrieval code (ndp057.sas) written by the preparers to read and print any of the files containing the station ¹⁴C data.

```
*****
* THIS SAS PROGRAM READS AND PRINTS THE NYDAL *
* AND LOVSETH ASCII FILES CONTAINING          *
* GROUND-LEVEL C14 MEASUREMENTS               *
*****
;
data nydal;
infile 'nydal.asc' missover firstobs=17;
input labref $ 1-9 week 11-15 samdat $ 17-30 @34 delC14 7.1 @42 delC13 5.1
      flag13 $ 47 corC14 50-55 sigma 60-61;
proc print;
run;
```

The following is a listing of the SAS® data retrieval code (strat.sas) written by the preparers to read and print the file containing the stratospheric ¹⁴C data.

```
*****
* THIS SAS PROGRAM READS AND PRINTS THE FILE  *
* (STRAT.DAT) CONTAINING C14 MEASUREMENTS      *
* DERIVED FROM AIR SAMPLES COLLECTED ON       *
* NORWEGIAN AIR FORCE AIRCRAFT DURING 1965     *
*****
;
data strat;
infile 'strat.dat' missover firstobs=17;
input sample $ 1-6 alt 11-15 samdat 27-32 delC14 45-48 @53 delC13 5.1
      flag13 $ 58 corC14 63-66 sigma 75-76;
proc print;
run;
```

15. PARTIAL LISTING OF DATA FILES

The following presents a partial listing of one (nydal.asc) of the 15 data files that contain the station ^{14}C data. This file is identical in format to all data files containing the station ^{14}C data.

The first 25 lines of the file nydal.asc are as follows:

```
*****
* Carbon-14 Measurements in Atmospheric CO2 from Northern *
* and Southern Hemisphere Sites, 1962-1993                *
*                                                          *
* Authors: Reidar Nydal and Knut Lovseth                  *
*       Radiological Dating Laboratory                     *
*       The Norwegian Institute of Technology              *
*       N-7034 Trondheim NTH                              *
*       NORWAY                                             *
*                                                          *
* NDP057 (November 1996)                                  *
*****
```

| Lab Ref. | Week | Sampling Interval | delta C14 | delta C13 | corr C14 | sigma |
|----------|------|-------------------|--------------|--------------|-------------|-------|
| D-00001 | 10 | 630304-630311 | 477.0 | -26.5 | 479.4 | 12 |
| D-00002 | 11 | 630311-630318 | 494.0 | -23.1 | 486.4 | 7 |
| D-00004 | 13 | 630325-630401 | 562.0 | -24.2 | 557.4 | 11 |
| D-00006 | 15 | 630408-630415 | 583.0 | -24.2 | 557.4 | 11 |
| D-00008 | 17 | 630422-630429 | 580.0 | -25.0 | 578.4 | 11 |
| D-00010 | 19 | 630506-630513 | 653.0 | -23.7* | 647.4 | 9 |
| D-00012 | 21 | 630520-630527 | 661.0 | -24.6 | 658.4 | 10 |
| D-00014 | 23 | 630603-630610 | 704.0 | -24.3 | 700.4 | 12 |
| D-00016 | 25 | 630617-630624 | 703.0 | -25.0 | 701.4 | 8 |

The last 25 lines of the file nydal.asc are as follows:

| | | | | | | |
|----------|-----|---------------|--------|--------|--------|----|
| TC-00109 | 576 | 740107-740114 | 435.0 | -23.7 | 428.1 | 10 |
| TC-00110 | 580 | 740204-740211 | 430.0 | -22.0 | 418.1 | 9 |
| TC-00111 | 584 | 740304-740311 | 427.0 | -22.4 | 416.1 | 9 |
| TC-00112 | 588 | 740401-740408 | 425.0 | -22.4 | 414.1 | 10 |
| TC-00113 | 595 | 740524-740601 | 410.0 | -23.2 | 402.1 | 10 |
| TC-00114 | 636 | 750303-750310 | 403.0 | -21.8 | 391.0 | 10 |
| TC-00115 | 680 | 760105-760112 | 396.0 | -21.6 | 382.9 | 11 |
| TC-00116 | 684 | 760202-760209 | 400.0 | -21.0 | 384.9 | 6 |
| TC-00117 | 688 | 760301-760308 | 392.0 | -18.6 | 370.9 | 8 |
| TC-00118 | 692 | 760329-760405 | 369.0 | -20.2 | 352.9 | 11 |
| TC-00119 | 696 | 760426-760503 | 357.0 | -24.2 | 350.9 | 9 |
| TC-00120 | 700 | 760524-760531 | 410.0 | -22.7 | 399.9 | 10 |
| V-00001 | 25 | 630619-630626 | 760.0 | -27.8 | 768.4 | 13 |
| V-00003 | 27 | 630701-630708 | 801.0 | -25.8 | 802.4 | 12 |
| V-00005 | 29 | 630715-630722 | 941.0 | -27.5 | 949.4 | 11 |
| V-00007 | 31 | 630729-630805 | 948.0 | -25.2 | 947.4 | 10 |
| V-00009 | 33 | 630812-630819 | 998.0 | -26.2 | 1001.4 | 11 |
| V-00011 | 35 | 630827-630903 | 1006.0 | -26.9 | 1012.4 | 13 |
| V-00013 | 37 | 630909-630916 | 994.0 | -26.4* | 997.4 | 12 |
| V-00015 | 39 | 630923-631001 | 960.0 | -28.1 | 970.4 | 10 |
| V-00017 | 41 | 631007-631015 | 964.0 | -26.4* | 969.4 | 11 |
| V-00019 | 43 | 631022-631029 | 949.0 | -22.8 | 938.4 | 11 |
| V-00055 | 79 | 640629-640706 | 878.0 | -26.8 | 883.3 | 11 |
| V-00057 | 81 | 640713-640720 | 973.0 | -26.4* | 976.3 | 11 |
| V-00059 | 83 | 640727-640803 | 919.0 | -26.4* | 922.3 | 11 |

The following is a complete listing of the data file (strat.dat) containing the stratospheric ^{14}C data.

```
*****
* Carbon-14 Measurements in Atmospheric CO2 from Northern *
* and Southern Hemisphere Sites, 1962-1993                *
*                                                           *
* Authors: Reidar Nydal and Knut Lovseth                  *
*           Radiological Dating Laboratory                  *
*           The Norwegian Institute of Technology           *
*           N-7034 Trondheim NTH                           *
*           NORWAY                                           *
*                                                           *
* NDP057 (November 1996)                                   *
*****
```

| Sample | Altitude (m) | Date of Collection (Year, Month, Day) | delta C14 | delta C13 | Corrected delta C14 | Sigma |
|--------|-----------------|--|--------------|--------------|------------------------|-------|
| AS-001 | 12600 | 650407 | 1459 | -8.5* | 1376 | 28 |
| AS-003 | 12000 | 650513 | 1660 | -8.5* | 1577 | 47 |
| AS-004 | 11900 | 650601 | 1140 | -8.5* | 1067 | 18 |
| AS-005 | 12000 | 650614 | 1718 | -8.5* | 1626 | 47 |
| AS-006 | 12000 | 650623 | 1669 | -8.1 | 1576 | 32 |
| AS-008 | 12000 | 650721 | 1437 | -8.5* | 1355 | 35 |
| AS-009 | 12000 | 650804 | 1116 | -11.2 | 1055 | 33 |
| AS-010 | 9000 | 650821 | 814 | -3.3 | 734 | 17 |
| AS-011 | 12000 | 650907 | 854 | -8.5* | 737 | 23 |
| AS-012 | 12000 | 650924 | 986 | -8.5* | 919 | 26 |
| AS-013 | 12000 | 651014 | 599 | -11.4 | 554 | 33 |

16. VERIFICATION OF DATA TRANSPORT

The data files contained in this NDP may be read with the FORTRAN 77 or SAS® data retrieval programs provided. To verify that the data have been correctly transported to their systems, users should generate some or all of the statistics presented in Tables 2 and 3. These tables present simple summary statistics for the file that contains data from all 14 sites (i.e., file **nydal.asc**) and the file that contains the stratospheric ¹⁴C measurements (i.e., file **strat.dat**). If the statistics generated by the user differ from those presented here, the data files may have been corrupted in transport.

These statistics are presented only as a tool to ensure proper reading of the data files. They are not to be construed as summaries of the data sets.

Table 2. Characteristics of numeric variables in the file containing carbon isotope data from all 14 sampling locations

| Variable | Number of observations | Mean | Standard deviation | Minimum value | Maximum value |
|----------|------------------------|-------------|--------------------|---------------|---------------|
| WEEK | 1619 | 534.8097591 | 490.3078555 | -34 | 1591 |
| DELC14 | 1619 | 502.1426189 | 237.0226744 | 108.5 | 1036.0 |
| DELC13 | 1619 | -24.2290303 | 2.3573591 | -31.7 | -11.3 |
| CORC14 | 1619 | 496.9402718 | 236.6761277 | 110.4 | 1040.4 |
| SIGMA | 1619 | 9.1432983 | 3.1888204 | 3 | 42 |

Table 3. Characteristics of numeric variables in the file containing stratospheric carbon isotope data derived from air samples collected by aircraft during 1965

| Variable | Number of observations | Mean | Standard deviation | Minimum value | Maximum value |
|----------|------------------------|-----------|--------------------|---------------|---------------|
| ALT | 11 | 11772.73 | 938.1800564 | 9000 | 12600 |
| SAMDAT | 11 | 650722.64 | 188.6336517 | 650407 | 651014 |
| DELC14 | 11 | 1222.91 | 387.7791780 | 599 | 1718 |
| DELC13 | 11 | -8.50 | 2.0712315 | -11.4 | -3.3 |
| CORC14 | 11 | 1143.27 | 380.9330889 | 554 | 1626 |
| SIGMA | 11 | 30.82 | 9.9981817 | 17 | 47 |

APPENDIX A

FIGURES SHOWING TROPOSPHERIC AND STRATOSPHERIC ^{14}C MEASUREMENTS IN ATMOSPHERIC CO_2 FROM 14 NORTHERN AND SOUTHERN HEMISPHERE SITES




Fig. A-1. Corrected ^{14}C measurements from air samples collected at Kapp Linné, Fruholmen, Gråkallen, and Vassfjellet




Fig. A-2. Corrected ^{14}C measurements from air samples collected at Lindesnes, Norway; Santiago de Compostela, Spain; and Rehovot, Israel






Fig. A-3. Corrected ^{14}C measurements from air samples collected at Izaña, Canary Islands; Mas Palomas, Grand Canary Island; Dakar, Senegal; Dakar-Fann, Senegal; and N'Djamena, Chad




Fig. A-4. Corrected ^{14}C measurements from air samples collected at Debre Zeit, Ethiopia; Fianarantsoa, Madagascar; and by aircraft flying in the low stratosphere above Trondheim, Norway

INTERNAL DISTRIBUTION

- | | |
|----------------------|--------------------------------------|
| 1. L. J. Allison | 15. J. M. Loar |
| 2. T. A. Boden | 16. W. M. Post |
| 3. A. L. Brenkert | 17. D. E. Reichle |
| 4. M. D. Burtis | 18. D. E. Shepherd |
| 5. M.-D. Cheng | 19. D. S. Shriner |
| 6. R. M. Cushman | 20. L. D. Voorhees |
| 7. M. P. Farrell | 21. T. G. Yow |
| 8. D. E. Fowler | 22. Central Research Library |
| 9. K. N. Gibson | 23-37. ESD Library |
| 10. S. G. Hildebrand | 38-39. Laboratory Records Department |
| 11. G. K. Jacobs | 40. Laboratory Records, ORNL—RC |
| 12. S. B. Jones | 41. ORNL Patent Section |
| 13. D. P. Kaiser | 42. Y-12 Technical Library |
| 14. P. Kanciruk | 43-243. CDIAC |

EXTERNAL DISTRIBUTION

- 244. Wallace C. Broecker, Columbia University, Lamont-Doherty Earth Observatory, Palisades, New York 10964
- 245. Michelle Broido, Acting Director, Environmental Sciences Division, Office of Health and Environmental Research, Department of Energy, ER-74, 19901 Germantown Road, Germantown, Maryland 20874-1290
- 246. Fred A. Donath, Director, Institute for Environmental Education, Geological Society of America, 1006 Las Posas, San Clemente, California 92673
- 247. Wanda Farrell, Environmental Sciences Division, Office of Health and Environmental Research, Department of Energy, ER-74, 19901 Germantown Road, Germantown, Maryland 20874-1290
- 248. Roger J. Francey, CSIRO, Division of Atmospheric Research, Private Bag #1, Mordialloc, Victoria 3195, Australia
- 249. Diana W. Freckman, Director, College of Natural Resources, 101 Natural Resources Building, Colorado State University, Fort Collins, Colorado 80523
- 250. C. D. Keeling, Scripps Institution of Oceanography, University of California, GRD 0220, 9500 Gilman Drive, 2314 Ritter Hall, La Jolla, California 92093
- 251. Bob M. Key, Princeton University, Geology Department, Princeton, New Jersey 08544
- 252. Alexander Kozyr, The University of Tennessee, Pellissippi Research Facility, 10521 Research Drive, Suite 100, Knoxville, TN 37932
- 253. Ingeborg Levin, University of Heidelberg, Institute für Umweltphysik, Im Neuenheimer Feld 366, 6900 Heidelberg, Germany
- 254. Knut Lövseth, Norwegian Institute of Technology, Radiological Dating Laboratory, Sem Saelandsvei 5, N-7034 Trondheim, Norway
- 255. Martin R. Manning, NIWA, P.O. Box 31-311, Gracefield Road, Lower Hutt, New Zealand
- 256-260. Reidar Nydal, Norwegian Institute of Technology, Radiological Dating Laboratory, Sem Saelandsvei 5, N-7034 Trondheim, Norway
- 261. Curtis Olsen, Environmental Sciences Division, Office of Health and Environmental Research, Department of Energy, ER-74, 19901 Germantown Road, Germantown, Maryland 20874-1290

262. Ingrid U. Olsson, Uppsala University, Department of Physics, Box 530, S-751, 21 Uppsala, Sweden
263. Bobbi Parra, Environmental Sciences Division, Office of Health and Environmental Research, Department of Energy, ER-74, 19901 Germantown Road, Germantown, Maryland 20874-1290
264. A. Patrinos, Associate Director, Office of Health and Environmental Research, Department of Energy, ER-74, 19901 Germantown Road, Germantown, Maryland 20874-1290
265. T.-H. Peng, NOAA/AOML, Ocean Chemistry Division, 4301 Rickenbacker Causeway, Miami, Florida 33149
266. G. S. Saylor, Director, Center for Environmental Biotechnology, The University of Tennessee, 1015 Research Drive, Suite 100, Knoxville, Tennessee 37932-2567
267. F. E. Sharples, Office of Science and Technology Policy, Environment Division, 443 Old Executive Office Building, 17th Pennsylvania Ave., NW, Washington, DC 20502
268. Minze Stuiver, University of Washington, Geological Sciences, C301 Health Sciences Building, Seattle, Washington 98105
269. Stuart C. Sutherland, Columbia University, Lamont-Doherty Earth Observatory, Climate/Environment/Ocean Division, Rt. 9W, Palisades, New York 10964
270. Taro Takahashi, Columbia University, Lamont-Doherty Earth Observatory, Climate/Environment/Ocean Division, Rt. 9W, Palisades, New York 10964
271. Pieter P. Tans, NOAA/CMDL, Maincode R/E/CG1, 325 Broadway, Boulder, Colorado 80303
272. Virgene Zumbunn, 216 Henry Hall, Princeton University, Princeton, New Jersey 08544
273. Office of Assistant Manager for Energy Research and Development, U.S. Department of Energy, Oak Ridge Operations, P. O. Box 2001, Oak Ridge, Tennessee 37831-8600
- 274-275. Office of Scientific and Technical Information, P. O. Box 62, Oak Ridge, Tennessee 37831